

# *Aesculus flava* 'Vestita'

## Vestia Yellow buckeye



### SEASONAL COLOURS



### TYPES OF PLANTING

**Tree types:** standard trees, multi-stemmed trees, shade trees | **Topiary on stem:** multi-stem umbrella

### USE

**Location:** street, avenue, square, park, central reservation, large garden, cemetery, traffic areas, industrial zones, countryside | **Pavement:** none, open, sealed | **Planting concepts:** Eco planting, Landscape planting, shade-tolerant

### CHARACTERISTICS

**Crown shape:** rounded, wide pyramidal | **Crown structure:** dense | **Height:** 15 - 18 m | **Width:** 10 - 15 m | **Winter hardiness zone:** 6A - 8A

### ASPECTS

**Wind:** intolerant to wind | **Soil:** loess, sabulous clay, peaty, heavy clay, light clay, sand, loamy soil, all soils | **Nutrient level:** moderately rich in nutrients, rich in nutrients | **Soil moisture level:** dry, moist, wet | **Light requirements:** sun, partial shade, shade | **pH range:** acidic, neutral, alkaline | **Host plant/forage plant:** bees, butterflies, nectar value 5, pollen value 5 | **Extreme environments:** no infection with bleeding canker, no infestation by horse-chestnut leaf miner

### PLANTKENMERKEN

**Flowers:** panicle, striking, standing, big | **Flower colour:** pale yellow | **Flowering period:** May - June | **Leaf colour:** buds yellow, dark green, underside yellow-green | **Leaves:** deciduous, ovate, obovate, palmate, opposite, underside hairy, serrate, composite | **Autumn colour:** yellow | **Fruits:** striking, toxic, capsule, husk | **Fruit colour:** brown | **Bark colour:** grey | **Bark:** peeling, furrowed | **Twig colour:** pale brown | **Twigs:** hairy | **Root system:** moderately branched, fleshy roots, tap root, central root